


In the Claims:

1. **(currently amended)** A composition comprising:

A) a lubricant; and

B) a mixture of antioxidants, wherein said mixture is prepared by the partial condensation of an alkylated diphenylamine ~~selected from the group consisting of mono-, di-, and~~ comprising a tri-nonylated diphenylamine with an aldehyde or ketone in the presence of an acidic catalyst to yield at least one acridan of the general formula:



along with residual alkylated diphenylamine remaining after said partial condensation;
wherein:

R₁, R₂, R₃, and R₄ are independently selected from the group consisting of hydrogen and nonyl, provided that at least ~~one~~ three of R₁, R₂, R₃, and R₄ ~~[[is]]~~ are not hydrogen, and R₄, and R₅ are independently selected from the group consisting of C₁ to C₂₀ hydrocarbyl and hydrogen; wherein, at the termination of said condensation, residual alkylated diphenylamine is not separated from the acridan product and remains in said mixture with said antioxidants.

2-4. **(cancelled)**

5. **(previously presented)** The composition of claim 1 wherein the alkylated diphenylamine is condensed with a ketone.

6. **(original)** The composition of claim 5 wherein the ketone is acetone.


7. **(previously presented)** The composition of claim 1 wherein the composition further comprises at least one antioxidant in addition to that provided by the mixture of acridan and residual alkylated diphenylamine.

8. **(original)** The composition of claim 7 wherein the additional antioxidant is selected from the group consisting of amine antioxidants, hindered phenol antioxidants, and mixtures thereof.

9. **(original)** The composition of claim 8 wherein the hindered phenol antioxidant is selected from the group consisting of 2,4-dimethyl-6-octyl-phenol; 2,6-di-t-butyl-4-methyl phenol; 2,6-di-t-butyl-4-ethyl phenol; 2,6-di-t-butyl-4-n-butyl phenol; 2,2'-methylenebis(4-methyl-6-t-butyl phenol); 2,2'-methylenebis(4-ethyl-6-t-butyl-phenol); 2,4-dimethyl-6-t-butyl phenol; 4-hydroxymethyl-2,6-di-t-butyl phenol; n-octadecyl-beta(3,5-di-t-butyl-4-hydroxyphenyl)propionate; 2,6-dioctadecyl-4-methyl phenol; 2,4,6-trimethyl phenol; 2,4,6-triisopropyl phenol; 2,4,6-tri-t-butyl phenol; 2-t-butyl-4,6-dimethyl phenol; 2,6-methyl-4-didodecyl phenol; tris(3,5-di-t-butyl-4-hydroxy isocyanurate); tris(2-methyl-4-hydroxy-5-t-butylphenyl)butane; 3,5-di-t-butyl-4-hydroxy hydrocinnamate; octadecyl-3,5-di-t-butyl-4-hydroxy hydrocinnamate; tetrakis{methylene(3,5-di-t-butyl-4-hydroxyhydrocinnamate)}methane; 1,2-bis(3,5-di-t-butyl-4-hydroxyhydrocinnamoyl)hydrazine; 1,3,5-tris(3,5-di-t-butyl-4-hydroxybenzyl)-s-triazine-2,4,6 (1H,3H,5H)trione; 2,2'-oxamido bis-{ethyl-3-(3,5-di-t-butyl-4-hydroxyphenyl)}propionate; 1,3,5-tris(4-t-butyl-3-hydroxy-2,6-dimethylbenzyl)-s-triazine-2,4,6-(1H,3H,5H)trione; 1,3,5-trimethyl-2,4,6-tris(3,5-di-t-butyl-4-hydroxybenzyl)benzene; 3,5-di-t-butyl-4-hydroxyhydrocinnamic acid triester with 1,3,5-tris(2-hydroxyethyl)-5-triazine-2,4,6(1H,3H,5H)-trione; bis(3,3-bis(4-hydroxy-3-t-butylphenyl)butanoic acid)glycolester; tetrakis{methylene (3,5-di-t-butyl-4-hydroxyhydrocinnamate)}methane; 1,3,5-trimethyl-2,4,6-tris(3,5-di-t-butyl-4-hydroxybenzyl)benzene; and 3,5-di-t-butyl-4-hydroxy-hydrocinnamic acid C₇-C₉ branched alkyl ester.

10. **(cancelled)**

11. **(currently amended)** A method for reducing the susceptibility of a lubricant to oxidation comprising adding to said lubricant a mixture of antioxidants, wherein said mixture is prepared by the partial condensation of an alkylated diphenylamine with an aldehyde or ketone in the presence of an acidic catalyst to yield at least one acridan of the general formula:



along with residual alkylated diphenylamine remaining after said partial condensation;
wherein:

R₁, R₂, R₃, and R₄ are independently selected from the group consisting of hydrogen and nonyl, provided that at least ~~one~~three of R₁, R₂, R₃, and R₄ ~~[[is]]~~ are not hydrogen, and R₄, and R₅ are independently selected from the group consisting of C₁ to C₂₀ hydrocarbyl and hydrogen; wherein, at the termination of said condensation, residual alkylated diphenylamine is not separated from the acridan product and remains in said mixture with said antioxidants.

12. **(currently amended)** The method of claim 11 wherein the alkylated diphenylamine is ~~selected from the group consisting of~~ comprises mono-, di-, and tri-nonylated diphenylamine.

13. **(original)** The method of claim 11 wherein the alkylated diphenylamine is condensed with a ketone.

14. **(original)** The method of claim 13 wherein the ketone is acetone.

15. **(original)** The method of claim 12 wherein the alkylated diphenylamine is condensed with a ketone.

16. **(original)** The method of claim 15 wherein the ketone is acetone.

17. **(original)** The method of claim 11 wherein the composition further comprises at least one antioxidant in addition to that provided by the mixture.

18. **(original)** The method of claim 17 wherein the additional antioxidant is selected from the group consisting of amine antioxidants, hindered phenol antioxidants, and mixtures thereof.


19. **(original)** The method of claim 18 wherein the hindered phenol antioxidant is selected from the group consisting of 2,4-dimethyl-6-octyl-phenol; 2,6-di-t-butyl-4-methyl phenol; 2,6-di-t-butyl-4-ethyl phenol; 2,6-di-t-butyl-4-n-butyl phenol; 2,2'-methylenebis(4-methyl-6-t-butyl phenol); 2,2'-methylenebis(4-ethyl-6-t-butyl-phenol); 2,4-dimethyl-6-t-butyl phenol; 4-hydroxymethyl-2,6-di-t-butyl phenol; n-octadecyl-beta(3,5-di-t-butyl-4-hydroxyphenyl)propionate; 2,6-dioctadecyl-4-methyl phenol; 2,4,6-trimethyl phenol; 2,4,6-triisopropyl phenol; 2,4,6-tri-t-butyl phenol; 2-t-butyl-4,6-dimethyl phenol; 2,6-methyl-4-didodecyl phenol; tris(3,5-di-t-butyl-4-hydroxy isocyanurate; tris(2-methyl-4-hydroxy-5-t-butylphenyl)butane; 3,5-di-t-butyl-4-hydroxy hydrocinnamate; octadecyl-3,5-di-t-butyl-4-hydroxy hydrocinnamate; tetrakis{methylene(3,5-di-t-butyl-4-hydroxy-hydrocinnamate)}methane; 1,2-bis(3,5-di-t-butyl-4-hydroxyhydrocinnamoyl)hydrazine; 1,3,5-tris(3,5-di-t-butyl-4-hydroxybenzyl)-s-triazine-2,4,6 (1H,3H,5H)trione; 2,2'-oxamido bis-{ethyl-3-(3,5-di-t-butyl-4-hydroxyphen-yl)}propionate; 1,3,5-tris(4-t-butyl-3-hydroxy-2,6-dimethylbenzyl)-s-triaz-ine-2,4,6-(1H,3H,5H)trione; 1,3,5-trimethyl-2,4,6-tris(3,5-di-t-butyl-4-hydroxybenzyl)benzene; 3,5-di-t-butyl-4-hydroxyhydrocinnamic acid triester with 1,3,5-tris(2-hydroxyethyl)-5-triazine-2,4,6(1H,3H,5H)-trione; bis(3,3-bis(4-hydroxy-3-t-butylphenyl)butanoic acid)glycolester; tetrakis{methylene(3,5-di-t-butyl-4-hydroxy-hydrocinnamate)}methane; 1,3,5-trimethyl-2,4,6-tris(3,5-di-t-butyl-4-hydroxybenzyl)benzene; and 3,5-di-t-butyl-4-hydroxy-hydrocinnamic acid C₇-C₉ branched alkyl ester.

20-22. **(cancelled)**

23. **(new)** A composition according to claim 1 comprising:

A) a lubricant; and

B) a mixture of antioxidants, wherein said mixture is prepared by the partial condensation of an alkylated diphenylamine comprising mono-, di-, and tri-nonylated diphenylamine with an aldehyde or ketone in the presence of an acidic catalyst to yield at least one acridan of the general formula:



along with residual alkylated diphenylamine remaining after said partial condensation;
wherein:

R₁, R₂, R₃, and R₄ are independently selected from the group consisting of hydrogen and nonyl, provided that at least three of is R₁, R₂, R₃, and R₄ not hydrogen, and R₄, and R₅ are independently selected from the group consisting of C₁ to C₂₀ hydrocarbyl and hydrogen; wherein, at the termination of said condensation, residual alkylated diphenylamine is not separated from the acridan product and remains in said mixture with said antioxidants.